

Highlights from ATLAS and CMS measurements Brookhaven Forum 2021

Kostas Theofilatos on behalf of ATLAS and CMS










ATLAS and CMS are particle detectors situated 100m underground recording data from the Large Hadron Collider



The **LHC** produces* about:

4000 W[±] s / sec

 $1200 Z^0 s / sec$

17 tt s / sec

 $1 h^0 s / sec$

 $\sim 0.007 \, h^0 h^0 \, s \, / \, sec$ SM prediction

*Run II (20 Hz / nb)

+ quark-gluon plasma



MC simulation

key ingredient for interpreting the LHC data and for designing better (S/B) experimental analyses

entering a new phase 10 y after **NLO** revolution


first NNLO simulations available (e.g. Powheg MiNNLO):

color singlet first (e.g., V, h^o), recently colored particles (tt) added, EW corrections available from several tools like Sherpa or MG5_aMC@NLO 3.X

new **PS** beyond **LL** accuracy, devised with weighting techniques allow internal variations avoiding adhoc comparisons, e.g., **Herwig** vs **Pythia** → new understanding of systematic uncertainties

the availability of the enormous amount of MC samples needed for the LHC analyses, comes with their own challenges on the computing side

development of **concurrent running**, first tests on **GPUs** e.g. **MG5_aMC@NLO**



S

ATLAS and CMS have produced ~100 new physics results in 2021, in this talk:


	ATLAS	CMS	ATLAS+CMS
SMP	2	3	5
ТОР	2	1	3
HIG	4	4	8
ВРН	1	1	2
HI	1	1	2
TOTAL	11	10	20

ATLAS physics results

CMS physics results

jets, V, V+jets, VV


Z+jets high PT



leading jet kinematics nicely captured from NLO MCs, inclusive P_T(II) is tricker (integrates all additional jets)

Z+jets high PT


integrated cross sections are also provided (in good agreement with the SM predictions)



$\Gamma(Z \rightarrow inv)$

1st measurement of the Z invisible width @ hadron collider


$$\Gamma(Z \to \nu \bar{\nu}) = \frac{\sigma(Z + jets)\mathcal{B}(Z \to \nu \bar{\nu})}{\sigma(Z + jets)\mathcal{B}(Z \to \ell \ell)} \Gamma(Z \to \ell \ell) \qquad \bullet \qquad \text{input from LEP}$$



 $\Gamma_{Z \to vv}$ is extracted as a scaling parameter on the $Z \to vv$ process, relative to $Z \to II$


$\Gamma(Z \rightarrow inv)$

major systematic uncertainties [1.5-2%] from lepton selection efficiencies and JES



competitive with the *direct* measurement @ LEP


WWW



WWW final state sensitive to triple & quartic couplings

SM expectation ~505 fb with pp→Wh→WWW* accounting for more than the half of the total cross section


$$\sigma(pp \rightarrow WWW) = 850 \pm 100 \text{ (stat.)} \pm 80 \text{ (syst.) fb}$$



first observation, 8.2σ significance

more on VV, see Darien Woods' talk


Inclusive Jets @ 13 TeV



CMS jet data together with the HERA DIS measurements and CMS top pair production cross section are used to constrain **PDFs**, a_s and m_{top} simultaneously

Inclusive Jets @ 13 TeV


Relative uncertainties in the gluon distribution




$$\alpha_{\rm S}(m_{\rm Z}) = 0.1177 \pm 0.0014 ({\rm fit}) \pm 0.0022 ({\rm model~and~param.}),$$
 $m_{\rm t}^{\rm pole} = 170.2 \pm 0.6 ({\rm fit}) \pm 0.1 ({\rm model~and~param.}) \, {\rm GeV}.$


Although jet data have no direct sensitivity to m_{top} , their influence on the gluon PDF and a_s can be reflected in m_{top}


Inclusive Jets @ 5 TeV





comparing with NLO, NNLO & different set of scales/PDFs







measured data are available (Rivet & HepData) for global fits


top

tt (all hadronic)






this analysis is particularly sensitive to 4-quark EFT operators and sets limits on a range of them




tt (all hadronic)

normalized differential cross sections




 $p_T(tt)$ sensitive to extra radiation, Powheg+Py8 with more ISR describes better the data compared to the nominal Powheg+Py8 m(tt) > 1 TeV, all predictions have similar level of agreement with the data


ttγ (dileptonic)



 $N_{Y} = 1$, $N_{bjet} \ge 1$, $N_{lep} = 2$, $m_{II} > 20$ GeV fiducial phase space

mtop






direct

from cross sections

interpreting m_{top}

in direct measurements, data are compared to multiple MCs assuming different values of m_t^{MC} , but m_t^{MC} is not trivially connected to m_t^{pole}

a new ATLAS analysis connects m_t^{MC} with a well defined pQCD calculation at NLL, $m_t^{MSR}(R = 1 \text{GeV})$ which is numerically close to m_t^{pole} , by calibrating the prediction to different templates of Powheg+Py8 MC




$$m_t^{\text{MC}} = m_t^{\text{MSR}} (1 \text{ GeV}) + 80_{-410}^{+350} \text{ MeV}$$

higgs

Higgs, so far


observation of all main:

production modes ggF, qqH, VH, ttH
& decay modes γγ, ZZ, WW, ττ, bb and evidence for μμ



within uncertainties, all measurements consistent (so far) with the hypothesis that the **H** boson is a **CP-even fundamental scalar**

H → tau tau (differential)




First differential measurement using H → TT mode


bins of $P_T(H)$, N_{jets} , $P_T(j_1)$

at high p_T, competitive precision wrt other final states e.g., ZZ, WW


H → tau tau (STXS)




contrary to the fiducial cross section measurements here the xs is measured per production mode in predefined mutually exclusive bins, allowing the combination with other decay modes



H → tau tau (STXS)



global signal strength





µqqH statistically limited

$$\sigma \mathcal{B}(H \to \tau \tau)$$
 (fb)

	•	,
Process	Measured	SM Prediction
Inclusive	2960^{+394}_{-370}	3422^{+172}_{-172}
ggH	3060^{+592}_{-552}	3051^{+160}_{-160}
qqH	$221^{+75.3}_{-73.3}$	$329^{+9.67}_{-9.67}$

H → tau tau (STXS)






total cross section


$$\sigma_{H\to\tau\tau} = 2.9 \pm 0.21(\text{stat})^{+0.37}_{-0.32}(\text{sys}) \text{ pb}$$

consistent with the CMS measurement $2960^{+394}_{-370}\,\mathrm{fb}$

ATLAS-CONF-2020-026 CMS HIG-19-015




$H \rightarrow \gamma \gamma (STXS)$



27 regions measured in total

*bins are merged in 17 regions for this plot, (minimal merging has instead 27 regions)

$H \rightarrow ZZ (STXS)$



clean and statistically limited, will benefit a lot with more data

ATLAS Preliminary Total \sqrt{s} = 13 TeV, 36.1 - 139 fb⁻¹ Stat. $m_{H} = 125.09 \text{ GeV}$ Syst. $p_{_{SM}} = 79\%$ SM Total Stat. Syst. + 0.08 + 0.07 ggF $\gamma\gamma$ 1.02 -0.11 (- 0.08 - 0.07 + 0.10 +0.11+ 0.04 0.95 ggF ZZ -0.11 0.10 -0.03+0.13+ 0.12 ggF WW 1.13 (- 0.06 , -0.10 A -0.12+ 0.15 +0.28+ 0.23 ggF ττ 0.87 -0.25 - 0.15 - 0.20 +0.91+ 0.77 + 0.49ggF+ttH μμ 0.52 -0.38-0.88-0.79+0.27 + 0.21 + 0.17 VBF γγ 1.47 -0.24- 0.20 - 0.14 +0.51 + 0.50 + 0.11 **VBF ZZ** 1.31 -0.420.42 , -0.06 +0.19+ 0.15 + 0.11 **VBF WW** 1.09 (-0.14 , -0.10 -0.17+ 0.14 + 0.15 +0.20VBF ττ 0.99 -0.14 , -0.12 -0.18+0.38+ 0.31 + 0.21 VBF+qqF bb 0.98 -0.36-0.33-0.15+ 1.32 +1.34+ 0.20 VBF+VH μμ 2.33 - 0.23 \ _{-1.24} , + 0.32 + 0.10 +0.33**VH** γγ 1.33 (-0.30)-0.08-0.31+ 1.14 + 0.24 +1.17 VH ZZ 1.51 -0.940.93 , -0.16 + 0.49 VH ττ 0.98 0.49 -0.57+0.28+ 0.19 + 0.20 WH bb 1.04 -0.26-0.19-0.18+0.24+ 0.17 + 0.17 ZH bb 1.00 -0.22-0.17 , -0.14+0.27+ 0.26 + 0.08 ttH+tH γγ 0.93 -0.06-0.25-0.24+ 0.44 +0.65+ 0.48 ttH+tH WW 1.64 (-0.43, -0.43)-0.61 + 1.69 + 1.65 ttH+tH ZZ 1.69 -1.10- 1.09 - 0.16 +0.86 +0.66ttH+tH ττ 1.39 -0.62-0.76-0.44+ 0.20 +0.34+0.28ttH+tH bb 0.35 -0.33 , - 0.27 -2 2 0 6 4 $\sigma \times B$ normalised to SM

ATLAS comb

ATLAS-CONF-2021-053

likelihood level combination of all decay modes

total signal strength

$$\mu = 1.057^{+0.064}_{-0.062}$$

the new ATLAS combination has **decreased uncertainty** wrt the previous one by ~10%

more about Higgs measurements in Giulia Di Gregorio's talk


beauty

precision charmed beauty


$$R_{D_s^{(*)+}/\pi^+} = \frac{\mathcal{B}(B_c^+ \to J/\psi D_s^{(*)+})}{\mathcal{B}(B_c^+ \to J/\psi \pi^+)}$$

 J/ψ used to trigger the decay of interest $B_c^+ \rightarrow J/\psi$ $D_s^{(*)+}$

 $B_c^+ \rightarrow J/\psi \pi^+$ as reference





All the ratios of branching fractions are well described by the theoretical predictions



exceeds the precision of all previous measurements

new kid on the block







 $\Xi_{\rm b}(6100)^{-}$

an excited beautiful strange baryon

its existence will help distinguishing among different models predicting the properties of the excited Ξ_b states







Kostas Theofilatos $W(\Xi_b \pi^* \pi^*) = W(\Xi_b) + W_{\Xi_b} \text{ [MeV]}$

heavy ions

Z+jet




the "medium" is transparent for the colorless Z⁰ but not for its hadronic recoil




p_T(Z⁰) is unaffected by the medium and informs us about the scale of the hadronic recoil

Z+jet



ratio of particle yields between Pb+Pb and pp collisions (I_{AA})

Z+jet



azimuthal decorrelations and p_T^{trk} losses (not show here) consistent with the hypothesis that the "hard" parton loses energy (i.e., jet quenching) and/or the medium induces modification of the parton shower.

Summary


Interesting physics with **heavy ion data** and decays of **B-hadrons**, from the **two general purpose detectors** @ LHC.

In the realm of the **standard model, top & higgs** many new results in 2021, more precision is desirable.


For each ~10% improvement in precision, we have to make non trivial advancements in the analyses.


In all shown, the role of **accurate MCs** for designing better analyses (S/B) and for interpreting the results is indispensable.

BACKUP



$H \rightarrow \gamma \gamma (STXS)$





$H \rightarrow ZZ (STXS)$

